

MISS LAURA FERTITTA (Orcid ID : 0000-0003-3528-9804)

DR. SMAIL HADJ RABIA (Orcid ID : 0000-0001-6801-7106)

Article type : Letter to Editor

Title: Immunological and virological profile
of children with chilblain-like lesions and SARS-CoV-2

L. Fertitta, MD^{1,2}, A. Welfringer, MD^{1,2}, A. Ouedrani, MD^{3,4}, L. Polivka, MD-PhD^{1,2}, S. Chhun, PharmD-PhD^{3,4}, L. Chatenoud, MD-PhD^{3,4}, J. Fourgeaud, MD⁵, S. Hadj-Rabia, MD-PhD^{1,2,6}, S. Temmam, PhD⁷, M. Eloit, PhD⁷, Sermet-Gaudelus, MD-PhD^{8,9,10}, C. Bodemer, MD-PhD^{1,2,6}

Affiliations:

1. Department of Dermatology, Université de Paris AP-HP-5, Hôpital Universitaire Necker-Enfants Malades, Paris, France.
2. Reference Center for Genodermatoses and Rare Skin Diseases (MAGEC), Université de Paris AP-HP-5, Hôpital Universitaire Necker-Enfants Malades, Paris, France.
3. Institut Necker-Enfants Malades, Université de Paris, Paris, France.
4. Laboratory of Immunology, Université de Paris AP-HP-5, Hôpital Universitaire Necker-Enfants Malades, Paris, France.
5. Laboratory of Virology, Université de Paris AP-HP-5, Hôpital Universitaire Necker-Enfants Malades, Paris, France.
6. INSERM U1163, Institut Imagine, Institut Imagine, Hôpital Universitaire Necker-Enfants Malades, Paris, France.
7. Pathogen Discovery Laboratory, Department of Virology, Institut Pasteur, Paris, France

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/JDV.16972](#)

This article is protected by copyright. All rights reserved

8. Department of Pneumology, Université de Paris AP-HP-5, Hôpital Universitaire Necker-Enfants Malades, Paris, France.
9. Reference Center Rare Diseases: cystic fibrosis
10. INSERM U1151, Institut Necker – Enfants Malades, Université de Paris, Paris, France.

Corresponding author: C. Bodemer, MD-PhD

Address: Department of Dermatology, Hôpital Universitaire Necker-Enfants Malades, 149 rue de Sèvres – 75743 PARIS CEDEX 15

Phone: + 33 1 44 49 43 87

Email: chirstine.bodemer@aphp.fr

Key words: COVID-19, SARS-CoV-2, chilblain-like lesions

Manuscript word count: 599, Tables count: 2, Figure count: 1

Funding sources: none

Funding and conflicts of interest: none

Dear Editor,

The link between SARS-CoV-2 and the reported cutaneous manifestations has not been established. We assessed a possible correlation between the paediatric dermatological manifestations and the biological investigations, using for the first time 3 different SARS-CoV-2 tests.

From April to June, 2020, minors presenting with skin manifestations and symptoms of COVID-19 themselves or any of first-degree relatives (i.e. fever, influenza-like, respiratory, Ear-Nose-Throat and/or digestive symptoms), were enrolled. Epidemiological and clinical information, description of households and biological results including 3 types of SARS-CoV-2 tests [nasal PCR (systemic symptoms within the past 48 hours), serology (IgG, techniques: Abbott ARCHITECT), and interferon- γ (IFN- γ)-ELISPOT-assay] were collected. IFN- γ -ELISPOT-assay, an early (since day 5) qualitative and quantitative analysis, evaluates specific memory T-cells by quantifying the IFN- γ production after a short-term stimulation with SARS-COV-2 peptide. At least one test among serology and IFN- γ -ELISPOT-assay was performed on patients with chilblains.

Thirty patients (20 boys, average = 9.5 years) representing 28 households were included. Thirty-seven symptomatic first-degree relatives were analyzed. In 23/30 patients (77%) and 14/17 (82%) of chilblains patients, COVID-19 was suspected in at least one first-degree relative and confirmed in 4 including 2 with chilblains.

Chilblains were reported in 17 patients with a large spectrum of severity (Figure 1). Lesions occurred before (n=2, average: 19 days), simultaneously (n=2, 12%) or after systemic manifestations (60%, average: 22 days). Spontaneous resolution was complete in an average of 27 days (10-50). Two patients relapsed in 15 and 45 days respectively. Other cutaneous manifestations occurred before (20%, average: 18 days), during (30%) or after systemic manifestations (50%, average: 25 days). Two patients, including 1 child, presented with a linear pattern of urticarial lesions, both also presented with chilblains (Figure 1).

Elevated CRP [average 14mg/L (0 to 200)] and/or increased inflammatory cytokines were noted in 11 children (37%) including 8/17 with chilblains (47%). Cytokine levels were increased in 58%, 50%, 40% and 33% of chilblains patients tested for: TNF- α (range 20-60pg/ml), IL-1 (range 7-280pg/ml), type 1 IFN (range 2-6UI/ml) and IL-6 (range 10-127pg/ml) respectively (Table 1). In chilblains, tests were performed in an average of 18 days (5-98) and 21 days (6-51) after skin lesions and systemic manifestations onset respectively.

The 3/3 nasal PCR were negative. Serology was positive in only 1/16 chilblains patient among the 26 patients tested. IFN- γ -ELISPOT-assay was negative in all the 10 chilblains patients tested. In children with chilblains, these tests were performed in an average of 45 days from lesions onset (5-82) and 56 days from systemic manifestations (35-89).

High levels of cytokines, mostly TNF- α , IL-1, type 1 IFN and IL-6 were noted in 47% of chilblains patients. Biological inflammation was not correlated with: 1/ time lapses from cutaneous or systemic symptoms to the blood test, 2/ severity of chilblains. A cytokine storm was described in adults with COVID-19 (8) and in the paediatric inflammatory multisystem syndrome temporally associated with SARS-CoV-2 infection (PIMS-TS): elevated CRP and IL-6 levels (9).

The peak of incidence of COVID-19 and the reported chilblains occurred simultaneously (1-7). In 28/68 reported patients presenting chilblains (5,7,10), serology was negative. Only 1/16 chilblains children serology was positive. Sensitivity of our technique varies from 100 to 85% in severe or mild symptomatic patients respectively. Moreover it is known to be positive after 19 and 30 days of evolution in 85% and 94% of the patients respectively. In all our patients, the COVID-19 was confirmed only once, using 3 different methods. Our result might reflect the estimated prevalence of seropositivity for SARS-CoV-2 in the general French population.

While epidemiological data, clinical manifestations and elevated cytokines level suggest an association with SARS-CoV-2, no evident link could have been made.

Acknowledgments

The patients in this manuscript have given written informed consent to the publication of their case details.

References

1. Dong Y, Mo X, Hu Y, Qi X, Jiang F, Jiang Z, et al. Epidemiology of COVID-19 Among Children in China. *Pediatrics*. 2020;145(6).
2. Recalcati S. Cutaneous manifestations in COVID-19: a first perspective. *J Eur Acad Dermatol Venereol JEADV*. 2020;34(5):212-3.
3. De Masson A, Bouaziz J-D, Sulimovic L, Cassius C, Jachiet M, Ionescu M-A, et al. Chilblains are a common cutaneous finding during the COVID-19 pandemic: a retrospective nationwide study from France. *J Am Acad Dermatol*. 2020. Epub.
4. Galván Casas C, Català A, Carretero Hernández G, Rodríguez-Jiménez P, Fernández Nieto D, Rodríguez-Villa Lario A, et al. Classification of the cutaneous manifestations of COVID-19: a rapid prospective nationwide consensus study in Spain with 375 cases. *Br J Dermatol*. 2020. Epub.
5. Andina D, Noguera-Morel L, Bascuas-Arribas M, Gaitero-Tristán J, Alonso-Cadenas JA, Escalada-Pellitero S, et al. Chilblains in children in the setting of COVID-19 pandemic. *Pediatr Dermatol*. 2020. Epub.
6. Colonna C, Monzani NA, Rocchi A, Gianotti R, Boggio F, Gelmetti C. Chilblain-like lesions in children following suspected COVID-19 infection. *Pediatr Dermatol*. 2020. Epub.
7. García-Lara G, Linares-González L, Ródenas-Herranz T, Ruiz-Villaverde R. Chilblain-like lesions in pediatrics dermatological outpatients during the COVID-19 outbreak. *Dermatol Ther*. 2020. Epub.
8. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet Lond Engl*. 15 2020;395(10223):497-506.

- Accepted Article
9. ECDC. Paediatric inflammatory multisystem syndrome and SARSCoV- 2 infection in children, 2020. <https://www.ecdc.europa.eu/sites/default/files/documents/covid-19-risk-assessment-paediatricinflammatory-multisystem-syndrome-15-May-2020.pdf>.
 10. El Hachem M, Diociaiuti A, Concato C, Carsetti R, Carnevale C, Ciofi Degli Atti M et al. A clinical, histopathological and laboratory study of 19 consecutive Italian paediatric patients with chilblain-like lesions: lights and shadows on the relationship with COVID-19 infection. J Eur Acad Dermatol Venereol JEADV. 2020. Epub.

Legends of the table and figure:

Table 1: Clinical and biological characteristics of the patients

Figure 1: Clinical picture of our paediatric series during the COVID-19 pandemic: chilblain-like lesions associated with livedo (A, B and C), spontaneous urticarial lesions with linear disposition (D and E).

Table 1: Clinical and biological characteristics of the patients

	Total patients, n=30 (%)	Chilblain-like patients, n=17 (%)
Mean age (extremes), sex ratio F:H	9.5 y (1.8-17.3), 0.5	11.2 y (1.8-17.3), 0.4
Household contact with a case of COVID-19:		
- Probable cases	23 (77)	14 (82)
- Confirmed cases: PCR / serology / both	2 (7) / 1 (3) / 1 (3)	2 (12) / 0 / 0
Households description:		
- Number of households	28	15
- Total of first degree relatives	79	13
Past medical history of patients:		
- Raynaud phenomenon, photosensitivity	0	0
- Auto-immune disease	0	0
- Inflammatory bowel disease	1 (3)	0
- Asthma / Atopic dermatitis	3 (10) / 2 (7)	3 (19) / 2 (12)
- Urticaria	1 (3)	1 (6)
- Obesity	0	0
Dermatological manifestations:		
- Chilblains: total / feet / hands /both	17 (57) / 14 (47) / 2 (7) / 1(3)	17 (100) / 14 (82) / 2 (12) / 1 (6)
- Eccrine hidradenitis	2 (7)	-
- Maculo-papular rash	8 (27)	-
- Urticaria	1 (3)	-
- Livedo	2 (7)	-
- Targetoid lesions	2 (7)	-
- Vascular / ecchymotic purpura	1 (3) / 1 (3)	-
- Erythema nodosum	1 (3)	-
- Mucosal manifestations	0	-
Average time of cutaneous complete remission	22 d (1-50)	27 d (10-50)
Symptoms:		
- Mean pruritus scale from 1 to 10	7 (1-10), n=11 (33)	6 (1-10), n=6 (62)
- Mean VAS pain scale from 1 to 10	6 (2-10), n=9 (27)	5 (3-8), n=4 (50)
Systemic manifestations:	n=20 (67)	n=10 (59)
- Fever	10 (33)	3 (18)
- Influenza-like symptoms	13 (43)	7 (41)
- Respiratory symptoms	10 (33)	6 (35)
- ENT symptoms / anosmia	10 (33) / 1 (3)	7 (41) / 0
- Digestive symptoms	7 (23)	3 (18)

- Articular symptoms	1 (3)	0
Mean time lapse from systemic symptoms to lesions:	n=20 (67)	n=10 (59)
- Systemic manifestations before	25 d (3-77), n=10 (50)	22 d (5-46), n=6 (60)
- Cutaneous manifestations before	18 d (2-30), n=4 (20)	19 d, n=2 (20)
- Simultaneous manifestations	0 d, n=6 (30)	0 d, n=2 (20)
Laboratory tests:	n=25 (80)	n=16 (94)
- Anemia (Hb<11g/dl)	1 (4)	0
- Hyperlymphocytosis (>5,2G/l)	2 (8)	0
- Neutrophilic hyperleukocytosis (>8G/l)	1 (4)	1 (6)
- Elevated liver enzymes (ALT, AST)	0	0
- Elevated creatinine	1 (4)	0
- Elevated CRP (>5mg/l), mean (extremes)	3 (12), 14 (0-200)	1 (6), 5 (0-49)
- Low PT (<70%)	2 (8)	1 (6)
- Elevated aPTT (ratio>1,2)	6 (24)	3 (19)
- Elevated fibrinogène (>3,5g/l) (n=15)	0	0
- Elevated D-dimer level (>500ng/ml) (n=15)	1 (7)	0
- Positive antinuclear antibodies, mean title (extremes), specificity (n=17)	14/17 (82), 264 (100-800), 0	11/14 (79), 263 (100-800), 0
- APLA syndrome (β2GP1, cardiolipin, lupus anticoagulant) (n=15)	1/15 (7)	1/12 (8)
- Positive C-ANCA, specificity (n=17)	4/17 (23), 0	2/14 (14), 0
- Complement anomalies: C3, C4, CH50 (n=9)	1 low CH50 at 66 (4)	1 low CH50 at 66 (7)
- Elevated cytokines serum concentrations:		
o IL1 (> 15pg/ml), mean (range)	6/14 (43), 105 (17-280)	6/12 (50), 104 (7-280)
o IL6 (>10pg/ml), mean (range)	5/15 (33), 59 (10-127)	4/12 (33), 62 (10-127)
o TNF-α (> 20pg/ml), mean (range)	7/15 (47), 33 (20-60)	7/12 (58), 31 (20-60)
o Type 1 IFN (α) (> 2UI/ml), mean (range)	4/12 (33), 4 (2-6)	4/10 (40), 4 (2-6)
Tests of SARS-CoV-2:		
- PCR positive	0/8	0/3
- IgG positive (Abbott ARCHITECT)	1/26 (4)	1/16 (6)
- ELISPOT	11/11 (100)	10/10 (100)
Mean duration of follow-up	34 d (8-72)	42 d (11-72)



jd_v_16972_f1.png